

Genomic selection in small holder systems: challenges and opportunities

Raphael Mrode, Julie Ojango, Okeyo Mwai

Animal Genetic Research for Africa (Biosciences for Farming in Africa),
Nairobi, 10-11 September 2015

Right click and open hyperlink to download pictures



Partner Logo

Introduction

- Single nucleotide polymorphism (SNP) is a DNA sequence variation occurring when a single nucleotide in the genome differs between paired chromosomes in an individual.
- Using the association between SNP and QTL for economic traits, genomic breeding values (GEBV) can be computed for animals
- Selection using GEBVs has been implemented for various livestock species and is termed genomic selection

Benefits of genomic selection


- Benefits of genomic selections have well been demonstrated in developed countries
 - Reduced generation interval
 - Accuracies of above 70% have been reported for young genomic proven bulls
 - Higher rates of genetic gains

UK data that demonstrated that genomics work

- Comparing 3001 bulls
 - April 2012 Young bull genomic indexes
 - December 2013 indexes for the same bulls, but now with daughter performance information

Genomic → Dtr proven (Apr-12 to Dec-13)

- Correlation = 0.88
- Average April 2012 = £100 PLI (rlb. 69%)
- Average Dec 2013 = £101 PLI (rlb. 81%) (+12% rlb increase)



Characteristics of genomic system in developed countries

- Genomic systems in developed countries are characterised
- With large reference populations
- Collaboration among countries
- Well defined phenotypes
- But mostly within pure breeds

Characteristics of genomic data in small holder systems

- Challenges:
 - Small data sets
 - Difficult to define good reference and validation populations
 - Little data on pure breeds
 - Mostly on cross-bred animals

Characteristics of genomic data in small holder systems

- Opportunities:
 - Genotypes on cross-bred animals
 - Explore novel methods adapted to the data structure
 - Collaboration – sharing data and methodologies applied

Opportunities offered by genomics in small holder systems

- In small holder systems data and pedigree recording poses a challenge.
- Quick wins from genotypic data includes
 - Reduces the need for accurate pedigree recording as genomic relationship can easily be computed
 - Parentage discovery using SNP data
 - Usage to determine the breed composition of cross-bred animals in the absence of pedigree.
 - Gives the opportunity to match different genotypes to different management systems
 - Future use in traceability of animal products

Demonstrate some of the challenges & opportunities in small holder genomic data

- Examine genomic prediction using the DGEA small holder data
- GBLUP with **G** matrix was formed
 - Using SNP alleles frequencies from all cross-breds for scaling
- Attempt was made to determine accuracy of genomic prediction from subsets of the population with different breed composition

The DGEA data

- Data consisted of 1038 cows genotyped with the 777K Illumina High density chip
- Cows from 5 random sites in dairy production areas in Kenya
- Test day milk records were available
- Yield deviations for milk yield were generated using a repeatability model

Classification of cows by breed composition.

Dairy%	Breedtype		Dairy%by-Breedtype
1 (0-20%)	Zebu	6	1
2 (0.33-35%)	Mixed + Zebu	6	2
3 (36-60%)	Ayr/Gue/Jer	1	3
3	Fri	2	4
3	Mixed + Zebu	6	5
4 (61-87.5%)	Ayr	1	6
4	Fri	2	7
4	Ayr/Fri	3	8
4	Gue/Fri	4	9
4	Ayr/Gue/Fri	5	10
4	Mixed + Zebu	6	11
5(>87.5%)	Ayr	1	12
5	Fri	2	13
5	Ayr/Fri	3	14
5	Gue/Fri	4	15
5	Gue/Fri/Ayr	5	16

Average relationship from G matrix using breed type

1 0.027

2 0.016 0.018

3 0.026 0.022 0.038

4 0.022 0.021 0.032 0.035

5 0.030 0.024 0.042 0.041 0.051

6 0.024 0.016 0.003 0.003 0.001 0.118

Maximum relationship from G matrix

1	0.590					
2	0.591	0.564				
3	0.608	0.586	0.602			
4	0.413	0.539	0.520	0.565		
5	0.242	0.308	0.564	0.561	0.484	
6	0.516	0.546	0.535	0.506	0.269	0.681

Accuracy of prediction for Zebu and Zebu+mixed crosses

mean & std	176	-0.6149	-0.5818	1.0750	0.2728
------------	-----	---------	---------	--------	--------

reg & corr	176	1.2563	0.3188		
------------	-----	--------	--------	--	--

mean & sd of diff	2	176	0.0332	1.0213	
-------------------	---	-----	--------	--------	--

Accuracy of prediction for cows with 61-87.5% Dairy

mean & std	448	0.0000	0.0276	1.3368	0.2386
reg & corr		1.9413	0.3465		
mean & sd of diff		0.0276	1.2740		

Accuracy of prediction for cows with greater than 87.5% Dairy

mean & std	297	0.3961	0.2868	1.5969	0.2200
reg & corr		2.9743	0.4098		
mean & sd of diff		-0.1093	1.5201		

Implications and opportunities of such accuracies

- Given data structure and size, accuracies of genomic prediction obtained are encouraging
- Can be used to identify extreme animals on performance
- Can be used to select team of young bulls to use as sires
- Reduction in generation interval with use of young bulls will result in increase rate of productivity

Implications and opportunities of such accuracies

- Need for more data.
- The common saying is that in the “era of genomics, the phenotype is king”
- Looking at smart and innovative digital ways to collect data in the proposed Africa dairy genetics gain project

Implications and opportunities of such accuracies

- Need for collaboration across countries. This has been key for application of genomics
- Only third of the 20,000 bulls in the reference pop for the German genomic system are home proven.
- In the UK, less than one third of 22,000 bulls in the reference pop are domestic bulls
- Across regional genomics might be necessary for application of genomics for small holder farmers
 - Possibly most exotics sires used in crossing breeding are similar across regions or countries
 - Policies that promote easy flow of data across country boundaries while maintaining data security and ownership will be needed

Implications and Opportunities of such accuracies

- Need to collaborate with developed countries where some of the sires of these cows could have been genotyped
 - Parentage discovery & determine breed composition
 - Genotype by environmental interaction if enough data
- Link small holder data to medium to large farms enabling applications of methods such as one step
- Genotypes on cross-breds offer a unique opportunity and approach to examining across breed genomics

Conclusions

- Genomics offers quick wins in small holder systems through use of genomic relationship matrix and parentage discovery reducing need for accurate pedigree.
- Given data structure for smaller , well adapted methodologies will be needed for genomic selection but initial results are encouraging
- Collaboration on across country or regional basis will be needed to ensure adequate data and best sires can be used across regions
- Genotyping of cross bred animals offers possibilities for further optimize cross breeding systems
- In general, genotypic data offers opportunities to model underling genetics for production and resilience and therefore provide ways to modify them even in wild life populations

better lives through livestock

ilri.org

ilri.org

better lives through livestock

ILRI is a member of the CGIAR Consortium

Box 30709, Nairobi 00100 Kenya

Phone +254 20 422 3000

Fax +254 20 4223001

Email ilri-kenya@cgiar.org

ILRI has offices in:

- Central America • East Africa
- South Asia • Southeast and East Asia
- Southern Africa • West Africa



The presentation has a Creative Commons licence. You are free to re-use or distribute this work, provided credit is given to ILRI.

CRP and CG logos

